AD-A268 841 TATION PAGE

Form Approved
OM8 No. 0704-0188

ed to average 1 hour per response, including the time for reviewing instructions, searching existing data sources inexing the collection of information. Send comments regarding this burden estimate or any other assoct of this rate. To wishington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson fice of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, OC 20503.

IT DATE

3. REPORT TYPE AND DATES COVERED Final Report 4/01/90-3/31/93

4. TITLE AND SUBTITLE

JSEP FELLOWSHIP: FRACTAL MOISE PROCESSES

FINAL REPORT

6. AUTHOR(S)

Steven B. Lowen, George W. Flynn, Malvin C. Teich

2305/AS

S. FUNDING NUMBERS

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Columbia Radiation Laboratory of Columbia University

Rm. 1001 Schapiro CEPSR

530 W. 120th St.

New York, NY 10027

8. PERFORMING ORGANIZATION REPORT NUMBER

the material

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U. S. Army Research Office

AFOSR

P. O. Box 12211

Research Triangle Park, NC 27709-2211

nÉ

10. SPONSORING/MONITORING
AGENCY REPORT NUMBER

AFOSR-90-0250

11. SUPPLEMENTARY NOTES

Ċ.

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION / AVAILABILITY STATEMENT

126. DISTRIBUTION CODE

Approved for public release; distribution unlimited

93-20271

STRACT (Maximum 200 words)

Three fractal stochastic processes were developed and defined. Their statistical properties were derived, and examples were provided. Fractal shot noise (FSN), which is formed when a homogeneous Poisson process is passed through a filter with a power-law decaying impulse response function, was defined. For certain parameters, FSN converges to fractional Gaussian noise, while for others it is a Levy-stable random process. The fractal shot-noise-driven Poisson process (FSNDP), formed when FSN becomes the rate function for a second Poisson process, was considered next. The FSNDP is a point process which has a power spectral density inversely proportional to frequency raised to an arbitrary power (between zero and unity) for certain ranges of parameters. Also developed, were two fractal renewal processes with power spectral densities similar to that of the FSNDP: a standard renewal process with arbitrary power between zero and unity, and a finite-valued, alternating renewal process where the power extends to two. Several applications of these three processes were examined in detail.

14.	SUL	NECT	TE	RM\$
-----	-----	------	----	------

)(值

15. NUMBER OF PAGES

- 6

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

18. SECURITY CLASSIFICATION

UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT

20. LIMITATION OF ABSTRACT

UNCLASSIFIED

UNCLASSIFIED

Standard Form 298 (Rev. 2-39)

JSEP FELLOWSHIP: FRACTAL NOISE PROCESSES

FINAL REPORT

GEORGE W. FLYNN AND MALVIN C. TEICH FOR STEVEN B. LOWEN

MAY 31, 1993

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

AWARD # AFOSR-90-0250

COLUMBIA RADIATION LABORATORY
OF COLUMBIA UNIVERSITY

APPROVED FOR PUBLIC RFLEASE

DISTRIBUTION UNLIMITED

Auceri	on for			
Disc	CARSI DI BARI DI BOSCODIO			
Justification				
Ву				
Dist ib ition /				
٨	vailability Codes			
Dist	Avail date (or Special			
A-1				

DTIC QUALITY INSPECTED 1

THE VIEWS, OPINIONS. AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHORS AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE AIR FORCE POSITION, POLICY OR DECISION. UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

Objectives

Major objectives were the definition, development, study, and application of several fractal noise processes.

Accomplishments

Three kinds of fractal noise processes were carefully defined, and their statistics exhaustively detailed. Numerous examples were considered.

Publications

- [1] S. B. Lowen and M. C. Teich, "Power-Law Shot Noise," *IEEE Trans. Inform. Theory* 36, 1302-1318 (1990).
- [2] S. B. Lowen and M. C. Teich, "Doubly Stochastic Poisson Point Process Driven by Fractal-Shot-Noise," Phys. Rev. A 43, 4192-4215 (1991).
- [3] S. B. Lowen and M. C. Teich, "Fractal Renewal Processes as a Model of Charge Transport in Amorphous Semiconductors," *Phys. Rev. B* **46**, 1816-1819 (1992).
- [4] S. B. Lowen and M. C. Teich, "Fractal Renewal Processes Generate 1/f Noise," Phys. Rev. E 47, 992-1001 (1993).
- [5] S. B. Lowen and M. C. Teich, "Fractal Renewal Processes," *IEEE Trans. Inform. Theory* 39, (Sept. 1993), in press.

Personnel

Steven Bradley Lowen received his Ph. D. degree in electrical engineering, in May, 1992. His thesis title was "Fractal Stochastic Processes." He is now an Associate Research Scientist with the Columbia University Department of

Electrical Engineering. Malvin C. Teich is Professor of Engineering Science in the Departments of Electrical Engineering and Applied Physics at Columbia University.

Technical Progress

They explored the behavior of fractal shot noise, for which the associated impulse response functions assume a decaying power-law form [1]. Fractal shot noise is a stationary continuous-time process that is fundamentally different from fractional Brownian motion. They obtained expressions for the moments, moment generating functions, amplitude probability density functions, autocorrelation functions, and power spectral densities for a variety of parameters of the process. For certain parameters the power spectral density exhibits 1/f-type behavior over a substantial range of frequencies, so that the process serves as a source of $1/f^{\alpha}$ shot noise for α in the range $0 < \alpha < 2$. For these parameters they presented the first-order amplitude probability density function. For other parameters the amplitude probability density function is a Lévy-stable random variable with dimension less than unity. This process then behaves as a fractal shot noise that does not converge to a Gaussian amplitude distribution as the driving rate increases without limit. They considered several physical processes that are well described by fractal shot noise in certain domains: 1/f shot noise. Cherenkov radiation from a random stream of charged particles, diffusion of randomly injected concentration packets, the electric field at the growing edge of a quantum wire, and the mass distribution of solid-particle aggregates.

Fractal shot noise may serve as the rate of a second Poisson process, since it is nonnegative by construction; the result is the fractal-shot-noise

driven doubly stochastic Poisson point process or FSNDP [2]. They explored the behavior of the FSNDP, for which the associated impulse response functions assume a decaying power-law form. For a variety of parameters of the process, they obtained expressions for the count number distribution and moments, Fano factor, normalized coincidence rate, power spectral density, and time probability densities. A number of these measures exhibit power-law dependencies, indicating fractal behavior. For certain parameters the power spectral density exhibits 1/f-type behavior over a substantial range of frequencies, so that the process serves as a $1/f^{\alpha}$ point process for α in the range $0 < \alpha < 2$. They considered two physical processes that are well described by FSNDP: Cherenkov radiation from a random stream of charged particles, and diffusion of randomly injected concentration packets.

 $1/f^D$ noise occurs in an impressive variety of physical systems, and numerous complex theories have been proposed to explain it. In addition to the FSNDP, they constructed two relatively simple renewal processes whose power spectral densities vary as $1/f^D$: 1) a standard renewal point process, with 0 < D < 1: and 2) a finite-valued alternating renewal process, with 0 < D < 2 [3-5]. The resulting event number statistics, coincidence rates, minimal coverings, and autocorrelation functions are shown also to follow power-law forms. These fractal characteristics derive from interevent-time probability density functions which themselves decay in a power-law fashion. A number of applications were considered: trapping and charge transport in amorphous semiconductors [3, 7], electronic burst noise, movement in systems with fractal boundaries, the digital generation of $1/f^D$ noise, and ionic currents in cell membranes.